

DESCRIPTIONS

OF

ONE HUNDRED NEW SPECIES

OF

HESPERIDÆ.

BY

W. C. HEWITSON.

PART II.

LONDON :

JOHN VAN VOORST, 1 PATERNOSTER ROW.

March 23rd, 1868.

apex: crossed obliquely from the inner margin by a broad black band indicative of the male.

UNDERSIDE as above, except that it is paler and irrorated throughout with grey.

Exp. $1\frac{5}{20}$ inch. *Hab.* Australia (Moreton Bay).

In the Collection of W. C. Hewitson.

Closely resembling in character the *Hesperia* of Europe, and *H. comma* and *Cernes* of the United States.

Genus CYCLOPIDES, *Hübner*.

CARTEROCEPHALUS, *Lederer*.

1. *Cyclopides Caicus*.

Alis fuscis: anticis maculis sex vitreis maculaque opaca albis, posticis macula magna discali alba: his infra macula discali maculis duabus tripartitis subcostalibus ochraceis: maculis duabus prope angulum analem albis.

UPPERSIDE dark brown. Anterior wing with six transparent white spots: two near the middle (one below the other), one beyond these, and three at the apex. Posterior wing with one large central silvery-white spot.

UNDERSIDE. Anterior wing as above, except that the costal and outer margins are rufous, the apex marked by two pale rufous spots, and that the two lowest white spots are united in one. Posterior wing rufous, with the central spot as above, two large ochreous trifold spots at the costal margin, two white spots near the anal angle, and a submarginal band of four small rufous spots.

Exp. $1\frac{5}{20}$ inch. *Hab.* Venezuela.

In the Collection of W. C. Hewitson.

Very near to *C. cypselus* of Felder.

2. Cyclopides Cænides.

Alis fuscis, ciliis rufis: anticis maculis sex vitreo-fulvis, posticis macula magna aurantiaco-fulva: his infra rufis immaculatis.

UPPERSIDE rufous-brown, darker towards the outer margin of the posterior wing: the fringe rufous. Anterior wing with six transparent orange spots: three longitudinally at the middle of the wing and three (very minute) near the apex. Posterior wing with a large oblong orange spot before the middle.

UNDERSIDE. Anterior wing as above, but paler near the margins. Posterior wing rufous, irrorated with paler colour.

Exp. 1 inch. *Hab.* Venezuela.

In the Collection of W. C. Hewitson.

Nearly allied to *C. cypselus* of Felder.

3. Cyclopides Carmides.

Alis fuscis: anticis maculis tribus vitreis in triangulum dispositis, posticis macula fasciaque aurantiacis: his infra macula argentea.

UPPERSIDE. Dark brown, with rufous hair at the base. Anterior wing tinted with purple towards the outer margin, with three central transparent spots placed in the form of a triangle. Posterior wing with an orange spot before the middle, and beyond it a band of three spots of the same colour.

UNDERSIDE rufous-brown. Anterior wing as above, except that part of the inner margin is pale. Posterior wing with a large silvery central spot.

Exp. $1\frac{8}{10}$ inch. *Hab.* Madagascar.

In the Collection of W. C. Hewitson.

4. Cyclopides argenteo-ornatus.

Alis fuscis: anticis maculis septem aurantiaco-flavis, posticis macula oblonga aurantiaca: his infra viridibus, maculis quatuor fasciaque argenteo-albis.

UPPERSIDE. Dark brown, the base clothed with rufous